

DETERMINATION OF PHOSPHOROUS LEACHING POTENTIALS ON SELECTED SOILS DERIVED FROM THREE PARENT MATERIALS IN AKWA IBOM STATE, NIGERIA

Umoh, F. O¹; Inyang A. A¹. and Akpan A. E. ²

1. Department of Soil Science, Akwa Ibom State University, Mkpato Enin, Nigeria.
, umohflorence@gmail.com
2. Department of Soil Science, Akwa Ibom State University, Mkpato Enin, Nigeria
, ednaakpan@yahoo.com

Abstract

This study was conducted to assess Phosphorus Leaching behavior of three (3) selected parent materials in Akwa Ibom State. The soils were formed from river alluvium, beach ridge sand and coastal plain sand. A treatment solution containing 0, 20, 40 and 80mg/l of P prepared from KH_2PO_4 were added to 20g of soil samples in duplicated cups, the upper cup perforated, mixed thoroughly and allowed to dry. The cups were carefully covered and allowed to incubate for 1, 7, 30, 60 and 90 days, respectively. The experiment was 3 x 4 factorial in completely Randomized Design with three (3) parent materials and four (4) rates of phosphorous at varying days. A total of 60 experimental units were generated for the incubation study. The soil samples were kept moist with distilled water at weekly interval throughout the duration of incubation. At the set days, the concentrations of P in the leachate were measured. The P concentrations with rates of P added were plotted against the days of incubations. The result shows that the leaching behavior of K in the parent materials varies under similar experimental conditions. Beach ridge sand ($52.1mg L^{-1}$) and coastal plain sand ($21.1mg L^{-1}$) with high sand content exhibits highest amount of leaching compared to river alluvium ($9.16mg L^{-1}$) with high clay and organic matter content. A higher P losses with high rates of P addition was observed and P decreased with time (days) of incubation in the studied soils. River alluvium has long lasting effect of added P with a strong correlation with clay while short lasting effect was observed in coastal plain sand and beach ridge sand with a weak relationship with sand.

Keywords: Phosphorous, leaching effect of P, parent material, incubation time, soil properties.

Introduction:

Phosphorous (P) is one of the essential nutrient elements required in large amount for growth and development of plants. When phosphorous is applied in the form of water soluble fertilizers, it can either be absorbed on the clay minerals, taken up by plants or leached out. The extent of each mechanism and component depends upon various factors including the nature of parent materials, soil physical and chemical properties, contact time, amount of P applied, rainfall or irrigation etc. (Alfaro *et al.*, 2004, Sharma and Sharma 2011). Leaching under Agricultural fields is regarded as the movement of nutrients from top soil down the profile beyond the root zone and potentially to

surface water by lateral movement or lower to under ground water (Fink *et al.*; 2016; Raphael *et al.* 2018). Fertilizers are applied at high rates to increase yield, yet little is known about the leaching potential of nutrients from soils into the surrounding and underground water. Barman *et al.*, (2012) reported that Basic cations are known to be quickly solubilized in the soil solution with high quantity lost to leaching which leads to poor plant growth and quality. Blerina *et al.*, (2016) carried out a potted experiment to evaluate phosphorous leaching using two fertilizers rates in the presence and absence of plants and observed a reduced nutrient leaching on the pots cultivated with Ryegrass plant compared to a

bare Pot with high leaching of P. Gikonyo *et al.*, (2010) reported that the amount of nutrient leached increase with increasing rate of fertilizer applied and total nutrient concentrations in the leachate varies with soils. Umoh *et al.*, (2018) carried out an experiment to assess the leaching behaviour of K on six soil types in Eastern Nigeria using column leach test. It was observed that soils with high sand content exhibits highest amount of leaching compared to soil with high clay content, a higher K losses with high rates of K addition was also observed and K decreased with time of incubation. The greater phosphorous loss to the surrounding leads to environmental pollution. Jarvie *et al.*, (2013); Armagan *et al.*, (2014) confirmed that Phosphate (PO_4) is associated with harmful algal blooms and eutrophication of lake and found to decrease water quality. Li *et al.*, (2018) reported that irrigation has more influence than fertilization of leaching, water quality and that the optimal irrigation combined with optimal fertilization was efficient in reducing the potential environmental risk cause by excessive fertilization. Also reported that pH, and concentration of P, K, Ca, Mg and Cu leach in soils was significantly decreased by irrigation under same fertilization condition. Fink *et al.*, (2016) observed a reduction in P mobility in sandy soils with small amount of Fe and Al ion present. Increasing costs of fertilizers, low yield of crops and concern with water quality have motivated interest to minimize losses through this medium and to improve the efficiency use of fertilizers. Information on the leaching potential of phosphorous in these soils is limited. The study is therefore aim at determining the risk of phosphorous leaching in these soils using suction cups experiment.

Materials and Methods

Description of the study area and soil sampling

The study area is in Akwa Ibom State, lies between Latitudes $4^{\circ}32'$ and $5^{\circ}33'N$ and longitudes $7^{\circ}25'$ and $8^{\circ}25'E$. The cultivated area falls in the rainfed zone dominated by maize, cassava, legume, palms. The dominant forest types in the area include; The saline water swamp, fresh water swamp forest and the rain forest. Characterized by heavy rainfall ranging from 2000mm in land to over 3500mm along the coast and temperature range between $26^{\circ}C$ to $28^{\circ}C$ within a year. (Peters, 1989). Three surface soil samples (0 to 20cm) were collected to represent different parent materials. They include: River Alluvium (Okon in Ikot Abasi), Coastal Plain Sand (Obio Akpa in Oruk Anam) and Beach ridge sand (Uta Ewa in Ikot Abasi). The soil samples were air-dried, crush to pass through 2mm size sieve and use for the analysis.

Soil Analysis

Parts of the soil samples sieved were used for the following analysis: Particle size distribution was analysed using hydrometer method Udo *et al.*, (2009), soil PH was determined in water 1:2.5 soil and water ratio in KCl using glass electrode as described by Udo *et al.*, (2009). Organic matter was determined by wet-oxidation method described by Udo *et al.*, (2009). Available phosphorous was determined by Udo *et al.*, (2009), after extraction by Bray P-I extractant. Exchangeable cations were extracted with neutral NH_4OAc . Calcium and Magnesium were determined in the extract by EDTA titration as described by Udo *et al.*, (2009). Potassium and Sodium by the use of flame photometer. Effective cation exchange

capacity (ECEC) was obtained by the summation of the exchangeable cations and exchangeable acidity as described by Udo *et al.*, (2009) and Base saturation % was calculated by sum of exchange cation x 100 ECEC.

Incubation Procedure

Twenty (20) grams of soil was weighed into a duplicated cup with a capacity of 23cm in diameter and 13cm in length and the upper cups perforated Ukpong *et al.*, (2014). A 20ml portion of the treatment solution containing 0, 20, 40, 80mg l⁻¹ prepared from (K H₂ PO₄) was added to each of the soil in the cups, mixed thoroughly for effective mixing of the P solution with the soils and allowed to dry. The cups were covered and allow to incubate for 1, 7, 30, 60 and 90 days respectively. The treatment combinations were then subjected into a Randomized Complete Block Design (RCBD) with a total of 30 sub samples. The soils were kept moist with 30ml distilled water at weekly intervals and covered for the duration of incubation. At the set days, the P in the leachate were determine using flame photometry to obtain available phosphorous.

Statistical Analysis

Analysis of Variance (ANOVA) were used to evaluate the differences among the soils. Pearson Correlation Coefficient were use to correlate the P obtained with some soil preperities

Result and Discussion

The physical and chemical properties of the soils are presented in Table Ia. The soil varied from sand to loamy sand. pH values in soils were low indicating moderately acid

condition. Available P in river alluvium was sufficient. The values falls within the critical level of 12 – 15mgkg L⁻¹ proposed by (Aduayi *et al.*, 2002). Organic matter content were low, falls below the critical level of < 20 and high 30gkg⁻¹. The order of abundance of exchangeable cation are (Ca > Mg > K > Na and were adequate in the soils (Okalebo *et al.*, 2002). The effective cation exchange capacity (ECEC) were low. The effective cation exchange capacity (ECEC) was high in all the soil with values remaining below 12 cmol/kg. The soils recorded high base saturation. This finding agreed with the work of Haulin and Beaton (2006) which stated that the release of nutrient by soils is influenced by the cations. The results obtained were significantly different from each parameter as indicated in Table Ib.

The amount of P obtained from the leachate is presented in Table 2. Beach ridge sand had the highest mean of P in the leachate while river alluvium had the least. The trend were as follows: beach ridge sand (52.1 mg L⁻¹) > coastal plain sand (21.1mg L⁻¹) > river alluvium (9.16 mg L⁻¹). The result also revealed higher extractable P with high rate of P application. This indicated that the higher rate of P applied resulted in higher losses of P. This seems obvious as more quantity of P was available for reaction with soil properties, which resulting in higher of P moving out of the soil column. This clearly indicates that not all applied P is absorbed on the exchange complex. This observation is strongly affirmed with the positive relationship with the extractable P (Table 3). This results are in agreement with those obtained by Sharma (2011); Ukpong *et al.*, (2014) who reported that light textured soils are more prone to leaching losses.

Table 1a: Physico-chemical Properties of the Soil

Soil Properties	River alluvium (Ikot Abasi)	Beach ridge sand (Ikot Abasi)	Coastal Plan sand (Obio-Akpa)
Sand } %	84.2	95.2	87..2
Silt }	6.8	1.4	3.8
Clay }	10.0	3.4	9.0
Texture	Loamy Sand	Sand	Sand
PH(H ₂ O)	5.4	5.0	5.1
Ec (dsm ⁻¹)	0.12	0.07	0.08
Organic Matter (gkg ⁻¹)	3.2	3.01	5.76
Total N gkg ⁻¹	0.09	0.08	0.14
Av. P gkg ⁻¹	15.74	11.32	13.72
Ex.Ca }	8.0	5.2	8.8
Mg }	2.1	1.6	2.6
Na }	0.07	0.06	0.08
K }	0.21	0.08	0.20
EA }	1.97	1.88	1.93
ECEC }	12.4	8.9	13.8
Base Saturation (%)	84.01	77.8	85.82

Table 1b: Physico-chemical Properties of the Soil

Significant at 5% probability level

PM	And	silt	clay	pH	EC	ORG	TN	AVP	Ca	Mg	Na	K	EA	ECEC	BS
BRS	95.14	1.44	3.42	5.0	0.07	3.01	0.08	11.32	5.2	1.6	0.06	0.08	1.88	8.92	77.8
CPS	87.14	3.86	9	5.1	0.08	5.76	0.14	13.72	8.8	2.6	0.08	0.2	1.93	13.61	85.82
RAS	83.14	6.86	10	5.4	0.12	3.2	0.09	15.74	8	2.1	0.07	0.21	1.97	12.35	84.05
LSD	6.11	2.72	3.55	0.21	0.11	1.54	0.03	2.21	1.89	0.50	0.01	0.07	0.05	2.43	4.21

Table 2: Amount of P extracted at different rates of P addition over different days from the leachate

Parent Material/Location	Rates of P added (mg/l)	Incubation Period Days					Mean
		Amount of P extracted from Leachate					
		1	7	30	60	90	
River alluvium (RSA)	0	10.30	4.67	10.70	10.70	1.38	
	20	18.70	6.67	11.30	11.30	1.06	
	40	13.00	8.00	12.00	12.30	2.80	
	80	12.00	15.30	17.30	18.70	2.70	
	0	13.8	8.66	12.8	8.57	1.99	9.16
Costal Plain Sand (CPS)	0	12.00	16.70	17.30	14.00	2.30	
	20	16.70	19.30	35.30	16.00	2.10	
	40	18.70	22.00	16.00	22.00	3.80	
	80	17.30	71.30	66.70	26.70	3.30	
			16.2	32.3	33.8	19.7	2.88
Beach ridge sand (BRS)	0	48.70	12.00	53.30	62.00	12.20	
	20	27.50	66.70	102.00	67.30	15.00	
	40	50.70	88.00	60.00	62.70	13.20	
	80	52.70	90.00	74.00	69.30	14.20	
			44.9	64.2	72.3	65.3	13.7

Effect of Leaching on Soil Parent Materials

The effect of leaching on parent material are shown on figure 1. The amount of available P in leachate at each time interval increased with increasing rate of P added, 0mg L^{-1} being the lowest and 80mg L^{-1} being the highest for the three soils studied. This reflected the fact that amount of P loss increase with increasing rate of addition. The rates of leaching in the soils showed no uniform pattern on incubation days (1 – 90 days). The P leaching in river alluvium soil was low inspite of high rates of addition. Beach ridge sands soils had the highest

leaching rate while river alluvium had the lowest leaching rates. The trend is as followed: Beach ridge sand (BRS) > coastal plain sand (CPS) > river alluvium (RAS). The high P losses in beach ridge sand soils could be attributed to the sandy nature of the soil, The low P losses in river alluvium soils could be due to high clay and organic matter content which has ability to retain applied P (Table 1). The result gives a clear evidence of a greater risk of P leaching from high rates of application. This is in agreement with the finding of Gurpal *et al.*, (2006) who observed high leaching rate of total P on five soils with four fertilization levels.




Fig 1: Effect of leaching behaviour on Soil Parent Materials

The Leaching Behavior of Phosphorous (P) in Soils.

The leaching rates of applied P in the study soils were obtained from the relationship between P concentration in the leachate at different rates of P added and at different incubation days and are presented in figures below, the letters A to D denote the rates of P added as A – Control (O) B – 20, C – 40 and D – 80mg kg⁻¹. The graphs are plots of the concentration of P in leachate and incubation days.

Fig 2: The plot of P. Concentration in leachate VS incubation days in river alluvium soil

It was observed from Figure 2 in alluvium soils that the rates of leaching increases with increasing rates of P added. The concentrations of P in the leachate were low and highest concentration within the rates added were obtained at 60 days of contact time and declined days after indicating long period of leaching effect of added P in alluvium soil, the long lasting effect of P in that soil could be attributed to high clay and organic matter content. (Table 1) and other soil properties with a positives relationship indicated in (Table 3). The results are in agreement with those obtained by *Ibia et al.*, (2008) who reported that phosphorous in solution exists as negatively charged phosphate ion, low leaching, extremely reactive and binds with aluminum, iron, Calcium when present in soil at high level. Umoh, *et al.*, (2014) also affirm the strong energy bonding of this monovalent cation (P) in the soil of this zone. This causes




the P to form new chemicals in the soil which held the nutrient tightly with the soil clay and organic matter.

Fig 3: Plot of P concentration in leachate VS incubation days in Coastal Plain Sand (CPS)

In coastal plain sand soils (figure 3), there was a gradual increase in the P leaching with rates of P added. The highest P losses was observed at 7th days of contact time and decreases at 30 days, indicating that P in coastal plain sand has short period of leaching effect of added P. This could be attributed to the Sandy nature of the parent material with large pore space Table 1. This showed that light texture soils are more prone to high leaching losses, as indicated in (Table 3) Which has a negative relationship with sand and fractional recovery. The results are in agreement with those obtained by Sharma and Sharma (2011); Umoh *et al.*, 2018 who reported higher leaching of nutrient in sandy loam soil.

Fig 4: Plot of P concentration in leachate VS incubation days in beach ridge sand (BRS)

In beach ridge sand soils (figure 4) a gradual increase in P losses with increase rates of P added in the soil was observed. The variation in days of incubation indicated an irregular pattern with high concentration as fitted in the figure. The highest P losses were observed at 30 day of incubation with the highest rate of P (80mg L⁻¹) added and declined at 60 days. This result revealed that the soils can give a significant yield reduction after 30 days of contact time. These findings could be attributed to the high sand content in that soil (Table 1) which also indicated a negative relationship with sand (Table 3). Sandy soils has weak interaction of nutrient in the soil solution (Kolachi and Jahali, 2007) which is



Correlations between soil chemical properties and fractional recovering in the leachate

Table 3 shows the correlations among the various soil parameters with recovering of P in the leachate, there was significant ($P \leq 0.05$) Positive correlation between the soil properties with the fractional

recovery (except) sand with a negative relationship indicated by the R^2 values in (Table 3). This trends shows that the nutrients were positively contributed to the leaching of phosphorous in the studied soils.

Table 3: Correlation Matrix between fractional recovery of added P soils with soil properties

	sand	silt	clay	Ph	EC	ORG	TN	AVP	Ca	Mg	Na	K	EA	ECEC	BS	PE_1	PE_7	PE_30	PE_60	PE_90
sand	1																			
silt	-0.968	1.000																		
clay	-0.982	0.903	1.000																	
Ph	-0.891	0.976	0.788	1.000																
EC	-0.408	0.167	0.575	-0.051	1.000															
ORG	-0.249	0.000	0.430	-0.217	0.986	1.000														
TN	-0.339	0.094	0.513	-0.125	0.997	0.996	1.000													
AVP	-0.990	0.994	0.945	0.946	0.276	0.111	0.204	1.000												
Ca	-0.854	0.698	0.938	0.525	0.823	0.716	0.779	0.773	1.000											
Mg	-0.655	0.446	0.787	0.240	0.957	0.895	0.933	0.542	0.952	1.000										
Na	-0.655	0.446	0.787	0.240	0.957	0.895	0.933	0.542	0.952	1.000	1.000									
K	-0.965	0.870	0.997	0.742	0.632	0.494	0.573	0.919	0.960	0.829	0.829	1.000								
EA	-0.992	0.992	0.950	0.941	0.290	0.126	0.218	1.000	0.782	0.554	0.554	0.925	1.000							
ECEC	-0.828	0.662	0.919	0.483	0.850	0.750	0.809	0.741	0.999	0.966	0.966	0.945	0.750	1.000						
BS	-0.855	0.699	0.938	0.527	0.822	0.715	0.778	0.774	1.000	0.952	0.952	0.961	0.783	0.999	1.000					
PE_1	-0.781	0.912	0.647	0.979	-0.252	-0.410	-0.322	0.861	0.342	0.039	0.039	0.591	0.853	0.296	0.344	1.000				
PE_7	-0.744	0.554	0.858	0.360	0.913	0.832	0.881	0.643	0.983	0.992	0.992	0.893	0.654	0.991	0.983	0.636	1.000			
PE_30	-0.842	0.950	0.724	0.995	-0.149	-0.312	-0.221	0.909	0.439	0.144	0.144	0.672	0.903	0.394	0.441	0.628	0.821	1.000		
PE_60	-0.792	0.919	0.661	0.983	-0.234	-0.393	-0.305	0.870	0.360	0.057	0.057	0.605	0.863	0.313	0.361	0.900	0.737	0.859	1.000	
PE_90	-0.804	0.927	0.676	0.986	-0.215	-0.376	-0.287	0.879	0.378	0.077	0.077	0.621	0.872	0.331	0.379	0.876	0.711	0.838	0.992	1.000

Conclusion

Physico-chemical properties of the soils shows that, soil were moderately acidic, light texture, salt free and some of the soils were low in nutrients. The concentration of P in leachate at different incubation period increase with increasing rate of P added and decrease with time (days) of Incubation. The soils of river alluvium had long lasting effect of added P and found to have low leaching potential, while soils from beach ridge sand and coastal plain sand had short lasting effect of added P and higher leaching potential. Therefore, for effective application and utilization of P, split application is recommended for these soils to reduced the risk of the nutrient leached out of the soil profile beyond the assess of the plant root or leached out to underground water and make farming practices more sustainable and community friendly.

References

Aduayi, E. A Chude, V. O., Adebuyi B. A and Olagiwola S. O (2002). Fertilizer Use management Practices for Crops in Nigeria. 3rd Ed. Produced by Federal Fertilizer Department, Federal Ministry of Agriculture and Rural Development, Abuja Nigeria Pp 23 – 73.

Alfaro, M. A. Jarvis, S. C. and Gregory, P. J (2004). Factors affecting potassium leaching in different soils. *Soil Use and Management Journal* 20(2): 182 – 189, ISSN 0266 - 0032

Amazan K. A, Faycal B, Bruna G, Alberto P. (2014). Managing N & P ro water bodies: Characteristion and solition to wards macro-Reginal Integrated Nutrient Management. JRC-Ispra, European Commission.

Barman, P. J., Kartha S. A, Gupta, S. and Pradhan, B. (2012). A study on leaching bahaviours of Na, Ca and K using column leach Test. World Academy of Science, Engineering and Technology. Pp 1034 – 2038.

Blerina, B. C, Evan R. and Adrian D. (2016). Evaluation of Nutrient leaching in Pots under Greenhouse condition, Albanian. *Journal of Agricultural Science* 15(1): 1- 7.

Conti L, Ceretta C. A, Ferreira P. A, Lorensini F, Brunetto G. (2015). Effects of Pig slurry application and crops of phosphorous content in soil and the chemical species in solution. *R Brusci solo* 39:774-787.

Fink J. R, India A. V, Bavaresco J, Barron V. Torrent J. Bayr. C. (2016). Adsorption and desorption of phosphorous in subtropical soils as affected by management system and mineralogy. *Soil Till Research*. 155:62-68.

Gikonyo, E. Zatharah, A. R, Hanafi, M. M and Anuar, R (2010). Phosphorous leaching in an acid tropical soil. “Recapitalized with Phosphate rock and triple superphosphate. *Scientific World Journal* W:1499 – 1508.

Gurpal, T., Sims, J. T and Haggard, B. (2006). Phosphorous Leaching in Soils: effect of soil type, soil phosphorous saturation and animal manures generated from modified diet. 18th World Congress of Soil Science Philadelphia, Pennsylvania, U.S.A 3:15-19.

Halvin, I. I and Beaton, J. D (2006). Soil Fertilizer. An introduction to Nutrient management prentice 7th Ed. Hall of India New Delhi Pp. 199 – 218.

- Ibia, T. O, Effiong G. S, Ogban, P. I and Obi, J. C (2008). Relationship between phosphorous forms and parent materials in soils of Southeastern Nigeria. *Acta Agromica Nigeria* 8(2):104-110.
- Jarvie, H. P Sharpley A. N, Spear B Kleinman, P. J. A (2013) Water quality remediation faces unprecedented challenges from legacy phosphorus “*Environ. Sci Technol.* 47, 8997- 8998.
- Kolahchi, Z. and Jalahi M. (2007). Effect of water quality on the leaching of Potassium from Sandy Soil. *Journal of Arid Environment* 68:624-639.
- Li. Y, Li. J, Geo. L, Tian. Y (2018). Irrigation has more influence than fertilization on leaching, water quality and the potential environmental risk in excessively fertilized vegetable soils. *Plus one Journal.pone* 13(19): eo20H570.doi:10.1371.
- Okalabo J. R, Gathua, K. W and Woomer, P. 1 (2002). Laboratory method of soil and plant analysis. A working manual 2nd ed. TSI 3F – (14T and Shek 12).
- Peters, S. W., Usoro, E. J., Obot, U. W and Okpon, S. N (1989). Akwa Ibom State Physical Background, Soils and Land Ecological Problems. Technical report of the task force on soils and land use. Government Print, Uyo. 602
- Raphael P. A, Ignacio H. S. Regina M. Q (2018). Mobility of phosphorous from organic and inorganic and organic source material in a sandy soil. *International Journal of Recycling of Organic Waste in Agriculture* 7:152-163.
- Schmieder, F. Bergstrom L., Riddle M. Gistafsson, J. P, Kirchmann, H. (2018). Phosphorous speciation in a long-term manure-amended soil profile – Evidence from wet chemical extracton, ³¹P – NMR and PK – edge XNES Spectroscopy. *Geoderma* 322:19-27.
- Sharma, V and Sharma, K. M (2011). Potassium Leaching from two texturally variable in Potato growing alluvial soils of North – *Western India Journal of the Indian Society of Soil Science* 59(4) 343-348.
- Udoh E, Ibia, O. T, Ogunwale, J. A, Ano, A. O and Esu, I. V (2009). *Manual of Soil, Plant and Water Analysis.* Sibon Book Ltd. Lagos.
- Ukpong I. M, Osodeke V. E and Akpan U. S (2014). Determination of phosphorous fixing capacities and residual values of soils of Akwa Ibom State Nigeria. *Merit Research Journals of Agricultural Science and Soil Science* 2(8):096-103.
- Umoh, F. O., Osedeke, V. E. Edem, I. D., Effiong, G. S., (2014). Application of Langmuir and Freundlich Models in Phosphate Sorption Studies in soil of contrasting parent materials in south-eastern Nigeria. *Open access library journal* 1:989-996
- Umoh, F. O, Osedeke, V. E and Akpan, U. S (2018). Assessment of Potassium Leaching Behaviours in selected soils of Southeastern Nigeria. *International journal of sustainable Agricultral Research.* 5(1):10-18