

INFLUENCE OF SEX ON RELATIONSHIP BETWEEN MORPHOMETRIC TRAIT MEASUREMENT AND CARCASS TRAITS IN BROILER CHICKEN RAISED IN HUMID TROPIC

Sam, I. M.

Department of Animal Science, Faculty of Agriculture, Akwa Ibom State University, Akwa Ibom State, Nigeria

Email: sidorenyin@yahoo.com

Abstract

This study was conducted to determine the influence of sex on relationship among carcass traits (carcass weight, breast, drumstick, thigh, wings) and morphometric traits (body weight, breast girth, keel length, shank length, thigh length, wing length and body length) in broiler chicken. More so, to evaluate correlated relationship between the morphometric traits and carcass traits. Sixty broiler birds were divided into two treatment groups (male and female) each group was replicated thrice with ten birds per replicate. The birds were reared in deep litter system till they attained 56 days of age; body weight and morphometric trait measurements were taken on day 56. Eighteen birds (nine from each group) were randomly selected, slaughtered and the carcass traits measured and recorded. The data obtained from these studies were subjected to Pearson correlation analysis. The correlated relationship among most carcass traits and morphometric traits were significantly positive ($p < 0.001$) with correlation coefficients ranging from 0.01-0.681 in males (M), females (F) and in combination of males and female (M+F). The correlation between morphometric traits and carcass traits showed that breast girth and live body weight were positively and significantly ($p < 0.001$) correlated with all carcass traits in all the sex groups. The observed significant relationship among the morphometric traits and carcass traits as well as between morphometric traits and carcass traits is an indication of the existence of high level of dependency amongst these traits, thus improvement in any one of these traits would elicit a correlated response in the other traits. More so, the positive and significant correlated relationship between carcass traits and breast girth indicate that breast girth can be used as indirect selection criteria for carcass traits in broiler chickens.

Key words: Broiler chicken, Correlation, Sex, Traits, Morphometric, Carcass

Introduction

Poultry production is regarded as a means of livelihood and a way of maintaining a level of economic independence. The carcass yield of broiler chicken is of primary concern to the producer and consumer. Thus, poultry farmers have to select broiler strains and sex that will

reach market weight at a reasonable age in order to maximize profit.

Information on correlation among the pre- and post-slaughter traits is quite important in poultry breeding. This is because knowing which of the pre-slaughter traits affects which of the post-slaughter traits enables the breeder to predict what kind of products to obtain at the end of

breeding programme (Mendes *et al.*, 2005; Mendes 2009; Yakubu *et al.*, 2009). Size and shape of various parts are the major determinants of the overall size and shape of the live bird or carcass. According to Pinto *et al.* (2006) body weight at various ages and carcass characteristics are examples of variables that can indicate the usefulness of chicken for commercial purposes.

Performance testing which forms the basis for breeding work is difficult to conduct in the case of slaughter value parameters (Yakubu *et al.*, 2009). Selection towards improving carcass characteristics requires reliable and easy to apply methods for estimating the performance and breeding value of poultry species (Kleczek *et al.*, 2007).

Broiler producers are now focusing on carcass yield especially with the payment of good prices for major components parts such as breast, thigh and drumstick. Some factors are known to affect carcass yield of broilers, such as sex, breed, age nutrition and management (Sam, 2006; Sam *et al.*, 2016; Havestein *et al.*, 2003, Horniakova and Abas 2009).

Several studies on correlation between carcass traits and linear body measurements showed high positive and significant correlations in poultry (Alphonsus *et al.*, 2012; Sam, 2006; Shafey 2003). The relationship between body weight and carcass traits was earlier studied by Musa *et al.* (2006), they reported significant positive correlation between live weight and carcass weight, breast muscle weight, leg muscle weight, heart weight, liver weight and abdominal fat weight, but significant negative correlation between live weight and abdominal fat percentage in Anak breed. Positive genetic correlations of body weight with abdominal fat weight and abdominal fat percentage was also reported by Deeb and Lamont (2002). In Anakbroiler strain, Ojedapo *et al.* (2008) found significant positive phenotypic correlations between live weight and carcass weight (0.95), shank weight (0.93) and breast muscle weight (0.97) but observed significant negative phenotypic correlation (-0.78) between live weight and carcass weight in Wadi Ross strain. The results obtained by Olawumi (2013) showed

that all the carcass traits (dressing weight, eviscerated weight, carcass weight, breast muscle weight, back weight, thigh weight and drumstick weight) measured were good indicators of live weight.

It is necessary to have a good knowledge of the relationship between linear body measurements and carcass traits in broiler chicken to enable the breeder or farmer to predict important carcass parts from linear body measurements. There is paucity of information on the relationship between linear body measurements and carcass traits in Nigeria, most studies conducted have been on relationship between linear body measurements and live body weight. Therefore, the objectives of this study are to establish correlated relationship between linear body measurements and carcass traits; and also to establish correlated relationship amongst the carcass traits in broiler chicken.

MATERIALS AND METHOD

Location and Climate

The experiment was conducted at poultry unit in the research and experimental unit of Akwa Ibom State University Poultry farm, Obio Akpa Campus in Oruk Anam Local Government Area of Akwa Ibom State. The area lies between latitude 4⁰30'N and longitudes 7⁰'E and 8⁰00'E of the Greenwich Meridian. The area is in the humid tropical region, characterized by two seasons, rainy and dry, which last respectively from March to November and from November to March. Other climatic parameters are; heavy rainfall of 2000-2500mm in the wet season, annual temperature range of 24⁰C-30⁰C, and relative humidity range of 75-79% (Wikipedia 2018).

Experimental Birds

A total of 60 day-old Abor acre broiler chicks were purchased from a reputable hatchery and were used for the experiment. The birds were raised together for seven weeks on deep litter pen, and were separated into sexes on the eight week using comb colour.

Feeding and Management

Prior to the arrival of the birds the pens were washed and disinfected. On arrival, the birds were received and put in the brooding house.

They were placed on intensive management system in a deep litter system. Their beddings were made up of dry wood shavings and high level of hygiene was maintained throughout the experimental period. The chicks were distributed into two groups of 30 birds each (males and females) and each group was replicated thrice and each replicate of 10 birds were housed in a pen measuring 2m x 2m. Commercial feed and water were provided *ad-libitum*. All necessary prophylactic medications and vaccines were also provided. The birds were weighed at the beginning of the experiment and weekly thereafter; live body weight was taken on the last day of (i.e day 56) of the experiment.

Slaughtering and Dissection Procedures

At the end of the experiment, 18 birds (9 males and 9 females) which were close to the mean body weight of the total birds in the two groups were used for carcass analysis. This was done after starving the birds overnight so as to obtain the true weight. The birds were weighed individually to obtain live body weight and linear body measurements were also done using a measuring tape (body weight, breast girth, keel length, shank length, thigh length, wing length and body length and results were recorded. The birds were slaughtered by cutting through the jugular vein with a sharp knife. The carcasses were scalded in hot water of about 80°C for a minute and the feathers were plucked manually. The carcasses were eviscerated by cutting through the vent and the viscera were removed. Weights were obtained for the whole carcass and major component parts vizly: carcass weight, breast, drumstick, thigh, wing and back.

Data Collection

Morphometric Traits Measurement

Prior to slaughtering, linear body measurements were obtained using a measuring tape. Data collected were:-

- i) Body weight: This was taken for individual birds measured in kg using measuring scale
- ii) Breast Girth: This was taken as the circumference of the breast around

the deepest region of the breast using measuring tape in cm.

- iii) Keel Length: It was taken as the length of the sternum in cm using measuring tape.
- iv) Shank length: was measured as the length of the tars-metatarsus from the hock joint to the metatarsal pad.
- v) Thigh Length: This was taken as the distance between hock joint and the pelvic joint using measuring tape in cm.
- vi) Wing Length: It is the distance between the tip of the phalanges and the coracoids-humorous joint using measuring tape in cm.
- vii) Body Length: It was taken as the distance between the tip of the beak and the longest toe within the nail as describe by (Yakubu *et al.*, 2009).

Carcass Traits Measurement

After slaughtering, data were also collected on carcass characteristics. Data collected were:

- i) Carcass Weight: It was taken as the hot eviscerated carcass without feet, head and abdominal fat using digital weighing scale.
- ii) Breast weight: measured as the weight of the breast
- iii) Drumstick: measured as weight of the drumstick
- iv) Thigh: measured as weight of the thigh
- v) Wing :measured as weight of the wing
- vi) Back: measured as weight of the back

The different parts were weighed using sensitive scale and were expressed in grams.

STATISTICAL ANALYSIS

Pearson's correlation coefficients were determined between live body weight, linear body measurements and carcass traits; and amongst the carcass traits using SPSS 2007 version.

Results and Discussion

Phenotypic Correlations amongst morphometric traits in broiler chicken

The phenotypic relationship among live body weight and carcass traits of male, female and combination of male and female (M+F) birds are shown in Table 1. In male birds, positive correlations were observed among morphometric traits ($r = 0.021$ to 0.681), most of which were significant except correlation between shank length and breast width ($r = -0.253$) and keel length and shank length ($r = -0.038$). In females, live body weight and morphometric traits were positively correlated and their correlation coefficient (r) ($r = 0.024$ to 0.438) except for correlation between breast width and breast girth ($r = -0.503$), breast girth and shank length ($r = -0.379$), breast girth and thigh length ($r = -0.297$), breast girth and

wing length ($r = -0.254$), keel length and thigh length ($r = -0.044$), keel length and body length ($r = -0.238$) which were negatively correlated. In both sexes (Male + Female combine), live body weight and morphometric trait measurements were positively correlated and their correlation coefficient (r) range from moderate to high (0.267 to 0.729). These findings are in agreement with the reports of Ige *et al.* (2007) who reported positive phenotypic correlations among live body weight and linear body measurement in local fowls. Sam, 2006 also had similar result working with Abor acre breed of broiler chicken. These results indicate that selection for one of these morphometric parts will lead to improvement in the other.

Table1: phenotypic correlations among morphometric traits in broiler chicken

	LW	BRW	BRG	KL	SL	TL	WL	BL
Males								
LW	1.00	0.345	0.400*	0.376*	0.057	0.465*	0.358	0.459*
BRW		1.00	0.092	0.625**	-0.253	0.376*	-0.043	0.025
BRG			1.00	0.295	0.326	0.200	0.681**	0.590**
KL				1.00	-0.038	0.312	0.087	0.422*
SL					1.00	0.021	0.286	0.207
TL						1.00	0.352	0.266
WL							1.00	0.609**
BL								1.00
Females								
LW	1.00	0.256	0.236	0.438*	0.238	0.024	0.285	0.238
BRW		1.00	-0.503	0.166	0.656**	0.459*	0.724**	0.223
BRG			1.00	0.015	-0.379	-0.297	-0.254	0.152

KL	1.00	0.132	-0.044	0.042	-0.238
SL		1.00	0.283	0.630**	0.227
TL			1.00	0.640**	0.492**
WL				1.00	0.542**
BL					1.00

Males + Females

LW	1.00	0.577**	0.349**	0.575**	0.346**	0.596**	0.596**	0.526**
BRW		1.00	-0.059	0.583**	0.469**	0.643**	0.604**	0.376**
BRG			1.00	0.218	-0.104	0.059	0.141	0.345**
KL				1.00	0.268*	0.410**	0.351**	0.267**
SL					1.00	0.419**	0.639**	0.372**
TL						1.00	0.729**	0.593**
WL							1.00	0.683**
BL								1.00

**P < 0.01, *P < 0.05, LW= Live weight, BRW= Breast width, BRG= Breast girth, KL=keel length, SL= shank length, TL= thigh length, WL= wing length, BL= back length

Phenotypic Correlations amongst Carcass Traits in Broiler Chicken

Correlation coefficients among carcass traits in males (M), females (F) and combination of male and female (M+F) are shown in Table 2. In male birds, positive correlations were observed in all the carcass traits except for breast weight and back weight (r = -0.351). In the female group there were positive correlation between carcass weight and breast weight (r = 0.692), carcass weight and thigh weight (r = 0.695) and breast and thigh weight (r = 0.832) but negative correlations were observed between some traits such as carcass weight and wing weight (r = -

0.077), drum stick and wing weight (r = -0.205), thigh weight and wing weight (r = -0.015), thigh weight and back weight (r = -0.061) also wing weight and back weight (r = -0.169). In the combined male and female (M+F) group there were positive high correlations between all the carcass traits. This therefore implies that selection for one carcass part will eventually lead to the improvement in other parts and can be used to predict the other. Similar result had earlier been reported in different strains of chickens (Ojedapo *et al*; 2008, Shafey *et al*; 2013, Sam 2006). Sam, 2006 reported significant positive correlations among linear body measurement in Abor acre broiler chicken.

Table 2: Phenotypic Correlations among carcass traits in broiler

Carcass	Drumstick	Breast	Thigh	Wing weight	Back weight
---------	-----------	--------	-------	-------------	-------------

	weight		weight		weight	
Male						
Carcass weight	1.00	0.534	0.728*	0.704*	0.808**	0.174
Drum stick		1.00	0.227	0.309	0.794*	0.777*
Breast weight			1.00	0.772*	0.653	-0.351
Thigh weight				1.00	0.595	-0.118
Wing weight					1.00	0.339
Back weight						1.00
FEMALES						
Carcass weight	1.00	0.624	0.692*	0.695*	-0.077	0.555
Drum stick		1.00	0.467	0.594	-0.205	0.475
Breast weight			1.00	0.832**	0.289	0.114
Thigh weight				1.00	-0.015	-0.061
Wing weight					1.00	-0.169
Back weight						1.00
Male + Female						
Carcass weight	1.00	0.816**	0.777**	0.820**	0.839**	0.496*
Drum stick		1.00	0.621**	0.869**	0.713**	0.243
Breast weight			1.00	0.673**	0.869**	0.706**
Thigh weight				1.00	0.735**	0.232
Wing weight					1.00	0.510*
Back weight						1.00

Phenotypic Relationship between morphometric traits and Carcass Traits in Broilers

The phenotypic correlation between morphometric traits and carcass traits in broiler chickens are presented in Table 3. The results indicate that in male broilers, carcass weight was positively correlated with all morphometric traits ($r=0.023-0.736$); breast weight had negative correlation with carcass traits except its relationship with breast girth which was positive and significantly correlated ($r=0.562$). The relationship between breast girth and all the carcass traits were positive, and highly significant with relationship with thigh weight being the highest ($r = 0.736$). In female broilers, all the morphometric traits were positively correlated with all carcass traits, with correlation coefficient ranging from 0.019 to 0.567, except the relationship between breast width and wing weight which was negative ($r = -0.311$).

When males and females were considered, it was observed that live body weight and breast girth had significant ($p<0.001$) positive correlations with all carcass traits measured, the correlation coefficient ranged from 0.344 - 0.623. The results of these study is in line with the reports of Raji *et al.* (2010), Musa *et al.* (2006), who reported significant positive correlations between live body weight and breast weight, wing and fat weights in their study. Ojedapo *et al.* (2008) also reported significant positive correlation between live body weight and carcass traits in Anka strain of broiler.

The high and significant correlation between breast girth and carcass traits in this study is in consonance with reports of several authors (Raji *et al.*, (2010) in chickens; Kleczel *et al.*, (2006) in Muscovy ducks and Vali *et al.*, (2005) in quails. The positive correlations between breast girth and carcass traits signifies pleiotropy effects of genes controlling these traits, thus suggesting that breast girth can be a good

indicator of carcass traits in poultry. The negative, low and non-significant correlation between some body measurements (keel length and breast width) and carcass traits indicates that these traits may not be considered as reliable predictors of carcass traits in broilers. Kleczek *et*

al., 2006 made similar observations in Muscovy ducks. Body measurement traits have been reported to play an important role in predicting the carcass weight of broiler chicken (Yakubu and Salako, 2009).

Table 3: Phenotypic correlation between morphometric traits and carcass yield traits in broiler chicken

TRAITS	CARSWT	BRSWT	DRMSTK	THGWT	WGWT
Male					
LW	0.029	0.211	0.536	0.023	0.411
BRW	-0.201	-0.560	0.398	-0.559	0.169
BRG	0.494	0.562*	0.313	0.736*	0.650*
KL	-0.197	-0.053	-0.783*	0.071	-0.416
Females					
LW	0.449	0.115	0.383	0.114	0.516
BRW	0.019	0.327	0.131	0.206	-0.311
BRG	0.449	0.049	0.392	0.034	0.317
KL	0.300	0.567	0.381	0.429	0.243
Males+ Females					
LW	0.344	0.481	0.486*	0.220	0.399
BRW	-0.628*	-0.783	-0.486*	0.599**	-0.443
BRG	0.438*	0.347	0.514*	0.625**	0.623**
KL	-0.472*	-0.538*	0.558*	0.285	-0.509*

**P < 0.01, *P < 0.05, LW= Live weight, BRW= Breast width, BRG= Breast girth, KL=keel length, SL= shank length, TL= thigh length, WL= wing length, BL= back length, CARSWT=Carcass weight, BRSWT= Breast weight, DRSTK=Drumstick, THGWT= Thigh weight, WGWT =Wing weight.

Conclusion

The study revealed moderate to high positive relationship among morphometric traits measurements and carcass traits suggesting that improvement in one of these morphometric traits will lead to a positive response in the carcass traits. The positive and significant correlated relationship between carcass traits and breast girth suggest that breast girth can be used as indirect selection criteria for carcass traits in broiler chickens.

References

Alphonsus, C., Sam, I. M., Opoola, E. and Ayigun, A. E. (2012). Correlated Relationship Amongst

Performance And Carcass Characteristics In Broiler Chickens Raised To Maturity. *Animal Productive Research Advances*, 8(1):50-53

Deeb, N. and Lamont, S. J. (2002) Genetic architecture of growth and body composition in unique chicken populations. *Heredity* 93: 107-118.

Havestein, G. B., Ferket, P. R and Quereshi, M. A. (2003). Carcass composition and yield of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poultry Sci.* 82(10): 1509-1518.

- Horniakova, E. and Abas, K. A. (2009). Influence of low levels of Protein and sex on carcass traits and nutrient content in broiler meats. *Slovak Journal of Animal Science* 42(20): 75-78.
- Ige, A. O., Salako, A. E., Ojedapo, L. O., Adedeji, T. A., Yakubu, A., Amao, S.R., Animasahun, A. O. and Amao, O. A. (2007). Prediction of body weight on the basis of body measurements in mature indigenous chickens in derived savannah zone of Nigeria. Proc. 32nd annual conference, Nigeria Society for Animal Production, 18-21 March, 2007, Calabar, Nigeria. Pp185-187.
- Kleczek, K., Wawro, W., Wawro, K. and Makowski (2007). Effect of body weights of day-old Muscovy ducklings on growths and carcass traits. *Arch Tierz Dummerst*, 50:204-213.
- Kleczek, K., Wawro, E., Wilkiewicz-Wawro, and W. Makowski (2006). Multiple regression Equations to estimate the content of breast muscles, meat, and fat in Muscovy ducks. *Poultry Science* 85(7):1318-1326.
- Mende, S., M. (2009). Multiple linear Regression models based on principal component scores to predict slaughter weight of broiler. *Arch. Cruflegelk* 73(2): 139-144.
- Mendes, M. A. Karabayir, I. E., and Savas, T. (2005) The relationship among pre and post slaughter traits of American Bronze, Turkey. *Arch. Tierz*, 48: 283-289.
- Musa, H. H., Chen, G. H., Cheng, J. H., Li, B. C. and Mekki, D. M. (2006). Study on carcass characteristics of chicken breeds raised under the intensive condition. *Int. J. Poul Sci.* 5(6): 530-533.
- Ojedapo, L. O., Akinoken, O., Adedesi, T. A., Olayeni, T. B., Ameen, S. A. and Amao, S. R. (2008) Effect of strain and carcass characteristics of three commercial broilers reared in deep Litter system in the Derived savannah area of Nigeria *World J. Agri Sci.* 4(4): 487- 491.
- Olawumi, S. O. (2013). Phenotypic correlations between live body weight and carcass traits in Arbor Acre breed of Broiler Chicken. *International Journal of Science and Nature*, 4(1):145-149
- Pinto, L. F., Dacker, C. M., Denrelo, M. R., Ledur, M. O and Contindlo, L. L. (2006). Principal component analysis applied to performance and carcass traits in chicken, *Animal Res* earch 55: 419-425.
- Raji, A. O., Igwebuike, J. U., Kwari, I. D. (2010). Regression models for estimating breast thigh and fat weight and yield of broilers from non- invasive body measurement. *Agriculture and Biology and Journal of North America*, 1(4): 469-475.
- Sam, I. M. (2006). Rearing methods and Growth curves for Body weight and major component parts of Broiler chicken raised to maturity. M. SC. Thesis. Ahmadu Bello University, Zaria.
- Sam, I. M., Ekpo, J. S., Ukpanah, U. A., Eyo, G. and Warrie, M. (2016). Relationship between Linear Body Measurement and Live Body Weight in West African Dwarf Goats in Obio Akpa *Journal of Biology, Agriculture and Healthcare*, Vol 6(16): 118-124.
- Shafey, T. M., Alodan, M. A., Hussein, E. O. and Al-Batshen H. A. (2013) the effect of sex on the Accuracy of predicting carcass composition of Ross Broiler chickens. *The Journal of Animal and Plant Science*, (23(4) 975-980.

SLUK-AK (1989). Soils and Land use studies. Government print office, Uyo, Akwalbom State. Soil Survey Staff 1994. Keys to soils taxonomy. Soil management support service (SMSS) technology. No. 19 p. 306.

SPSS.(2007). Scientific Package for Social Sciences.

Vali, N., Edriss, M. A. and Rahmani H. R. (2005). Genetic Parameters of Body and Some Carcass Traits in Two Quail Strains. *International Journal of Poultry Science*, 4:296-300.

Wikipedia (2018). The free Encyclopedia <http://en.wikipedia.org/wiki/Akwa-Ibom>.

Yakubu, A., K. L Wahor, K. O., and Agade, V. I. (2009). Using factor scores in multiple linear regression model of predicting the carcass weight of broiler chicken using body measurements. *Revista Cientifica Agricola*, 9(4): 963-967.

Yakubu A and Salako A E. 2009. Path coefficient analysis of body weight and morphological traits of Nigerian indigenous chickens. *Egyptian Poultry Science* **29**: 837-50.