

CLIMATE CHANGE, FOOD SECURITY, NATIONAL SECURITY AND ENVIRONMENTAL RESOURCES (GLOBAL ISSUES & LOCAL PERSPECTIVES)

CLIMATE CHANGE, FOOD SECURITY, NATIONAL SECURITY and ENVIRONMENTAL RESOURCES

GLOBAL ISSUES & LOCAL PERSPECTIVES

Edited by

Ahmed Makarfi

Ignatius Onimawo

Prince Mmom

Ani Nkang

Abdullahi Mustapha

Eteyen Nyong

PUBLISHED BY:

Society for Agriculture, Environmental Resources & Management (SAEREM)

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

First published 2024

SAEREM World

Nigeria

C 2023 Eteyen Nyong

Typeset in Times New Roman

All rights reserved. No part of this book may be reprinted or reproduced or utilized in any form or by any electronic, mechanical, or others means, now, known or hereafter invented including photocopying and recording or in any information storage or retrieved system, without permission in writing from the copyrights owners.

CLIMATE CHANGE, FOOD SECURITY, NATIONAL SECURITY AND ENVIRONMENTAL RESOURCES (GLOBAL ISSUES & LOCAL PERSPECTIVES)

ISBN 978-978-60709-9-5

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

Printed at: SAEREM World

TABLE OF CONTENTS

Preface

Editorial Note

Table of Contents

Acknowledgement

Dedication

Part one: CLIMATE CHANGE

Chapter 1:

The Concept of Technical Efficiency and Effects Climate Change on Palm Oil Processing

Eteyen Nyong

Chapter 2:

Enviromental Resource Policy: Forestry and Climate Change Challenges.

Bolaji, K.A , Kabir G.H and Arowolo O.V.

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

Chapter 3:

A Review of the Impact of Bush Burning on the Environment: Potential Effects on Soil Chemical Attributes

Chiroma, A. M.,^{1*} and Alhassan, A. B.,¹

Chapter 4

Effect of Climate Change on Income and Constraints of Periwinkle Harvesters in Nigeria

Eteyen Nyong

Chapter 5:

The Nexus between Climate Change and Agricultural Production in Nigeria

¹Ettah, O. I., ²Igiri, Juliana and ³Ettah, Goddy I.

Chapter 6:

Climate Change and Adaptation Management Practices in Crop and Animal Production.

Idris, Rakiya Kabir and Suleiman, Akilu

Part two: FOOD SECURITY

CHAPTER 7

Trend of Climate Change Variables: Food Security and Perception on Arable Crop Farmers in South-South Nigeria.

Eteyen Nyong

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

CHAPTER 8

Social Media Marketing Culture As an Innovation of Delivering Growth in Post-Covid-19 Era

Sadiq Mohammed Sanusi¹ and Ahmad Muhammad Makarfi²

CHAPTER 9

Digital Agricultural Marketing as A New Age Technologies in Post- Covid-19 Era

Sadiq Mohammed Sanusi¹ and Ahmad Muhammad Makarfi²

CHAPTER 10

Climate Change, Pollution and National Insecurity

Ogbanje, Elaigwu Christopher & Umar, Haruna Suleiman

CHAPTER 11

Insecurity: Impacts on Agro-Allied Industries and Food Production

Salami, Azeez Oyeniya

CHAPTER 12

**Evolution of Desert Encroachment Narratives and how it affects
Desertification Policy Implementation in Nigeria**

Abdullahi Umar; Abdullahi Adamu; Kabiru Shehu; Ismail Alfa Adamu and Sadiq Abdullahi

CHAPTER 13

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

Soil Conservation Management: Climate Change and Food Sufficiency

Eze, Kingsley Chijioké* Obasi, Nnenna Patrick and Inyang, Otoobong Anwanabasi.

CHAPTER 14

A Review of the Impact of Bush Burning on the Environment: Potential Effects on Soil Physical Attributes

Alhassan, A. B.,^{1*} and Chiroma, A. M.,¹

CHAPTER 15

Effect of Carbon Dioxide (CO₂) Emission on Rice Production in Nigeria

¹Ibrahim Mohammed Kebiru, ²Husseini Salihu, ¹Shaibu Ufedo Monday TURKEY

Chapter 16

Turkey Farming

Balogun, B.I PhD

Part three: NATIONAL SECURITY

Chapter 17

Anthropogenic Activities: Implications on the Population and Diversity of Fauna-Avifauna Species of old Oyo Forest

Adedoyin, S.O., Omifolaji, J.K., Jatto, S.O.S., Oluwagbemi, T., and Sale, F.A.

Chapter 18

Conservation of Forest Resources in Nigeria: Case Study of Indigenous Forest Food Plants Species

Okonkwo, H. O, Nsien, I. B., and Akomolede, L. A

Chapter 19

Duck Farming

Balogun, B.I PhD

Chapter 20

Poaching and Trade in Wildlife Products: A Global Perspective

Okonkwo, H. O, Nsien, I. B., and Akomolede, L. A.

Chapter 21

Peace Education and Critical Peace Education: Eradicating Violence and Promoting Peace in Nigerian Schools

Abdulganijy Aremu SULYMAN and Duze Daniel ALI

Chapter 22

Idealist Education and PEANism as Panaceas for Security Challenges in Nigeria

Abdulganijy Aremu SULYMAN and [Kassim A. OYEWUMI](#)

Part four: ENVIRONMENTAL RESOURCES

Chapter 23

Soil Conservation Management: Climate Change and Food Sufficiency

Lukumun Lekan, ADELAKUN

Chapter 24

Environmental Conservation: Food Production, Resource Management, Food Security, and Sustainability

Adeyemi Patrick OYEKAN

Chapter 25

Analysis of Green Leafy Vegetable Profitability and Risk Management among Women Marketers in Ekiti State, Nigeria

Ajibade, Y.E.*¹, Folayan, J.A.²., Akinyemi, M³., Ayeni, M.D⁴., Musa, V.H⁵., and Oni, S.O⁶.

Chapter 26

Environmental Communication: The Media and Climate Change Issues

Triumph-Aruchi Eteyen Nyong

Chapter 27

Ecotoxicology and Micro Bioindicators Assessment of Environmental Pollution

Mansur Abdul Mohammed

Chapter 28

Pigeon Farming

Balogun, B.I PhD

Preface

This book adopts an exegetical approach as well as a pedagogic model, making it attractive agriculture and environmental economics teachers, professional practitioners and scholars. It eschews pedantry and lays bars the issues in such clarity that conduces to learning. The book elaborates on contemporaneous climate change, food security, national security and environmental resources issues of global significance and at the same time, is mindful of local or national perspectives making it appealing both to international and national interests. The book explores the ways in which climate change, food security, national security and environmental resources issues are and should be presented to increase the public's stock of knowledge, increase awareness about burning issues and empower the scholars and public to engage in the participatory dialogue climate change, food security, national security and environmental resources necessary in policy making process that will stimulate increase in food production and environmental sustainability.

Climate Change, Food Security, National Security and Environmental resources: Global issues and Local Perspectives is organized in four parts. Part One deals with Climate Change with Six Chapters, Part Two is concerned with Food Security with Nine chapters, Part Three deals with National Security with Five Chapters, while Part Four pertains Environmental Resources, has Five Chapters.

Ahmed Makarfi / Eteyen Nyong

April 2024

CHAPTER 28

Pigeon Farming

Balogun, B.I PhD

Department of Agricultural Education, Federal University of Education, Zaria,
balogunib@gmail.com,

Introduction

The domestic or common pigeon (*Columbia livia domestica*) is an inclusive name for birds of the family *Columbidae*, a cosmopolitan group with about 300 species (NRC, 1991). Generally, larger species are called pigeon and smaller ones are known as doves, although there is no technical distinction. All pigeons trace back to a common ancestry, the blue rock pigeon. Pigeons have been man's closest associate for over 6000 years (Jin, *et al.* 2023). In the process of domestication they have branched off into varieties, fostered and developed by man for utility purposes (Didier, 2022; NRC, 1991). Pigeons are usually kept as show birds or for sport as racing birds and their well developed homing instinct, which means that they will fly back to their own roof space from wherever they are released, is well known. This feature has been used in the past as a means of communication with the message being borne in the pigeon's leg ring. Reuters, the international news agency has in the past transmitted its news via the humble pigeon. The birds have been known to travel for up to 700km. The other aspect of pigeon rearing is that of raising squabs for meat.

Pigeon are durable birds that can be raised with little effort and are able to survive harsh climates (Sule *et al.*, 2024; Arockia and Indianraj, 2023). They fend for themselves often ranging over many squares to locate seed and edible scraps. They have been raised for centuries especially in North America and Europe where they are produced as a delicacy for the market. But raising pigeons for food is not nearly as widespread as it could be: indeed, in modern times its potential has hardly been explored. Farmed pigeons are particularly promising as urban micro-livestock because they require little space and thrive in cities (NRC, 1991). Young pigeons (squabs) grow at a rapid rate. Their meat is finely textured with attractive flavour, and they are often used in place of game fowl. It is tender and easily digested, it commands premium market prices in many cases, the continuing demand is unfilled (NRC, 1991). According to Balogun (2009), pigeon meat has good protein content (19.8%), which is relatively comparable to duck (22.7%); local chicken (21.5%); quail

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

(28.3%); guinea fowl (26.4%) and exotic chicken (22.2%). It has low fat content of 1.3%. Pigeons are important commercial poultry in addition to being ornamental birds (Jin *et al.* 2023). In the year 2021, over 111million pairs of breeding pigeons were kept in stock and 1.6billion squabs were slaughtered for meat in China (Gao, 2022). According to Ji *et al.* (2022), the scale of pigeon business is expanding steadily in China with pigeon becoming the 4th largest poultry product after chicken, duck and geese. However, the squab sales has also increased over the years in the United States of America, Canada, Great Britain and France culminating into a drop in China's dominance of production from 80% of the total global production (Kokoszynski *et al.* 2000; Gao, 2022, Ji *et al.* 2022, Zhao *et al.* 2022). In Africa, the commercial production of pigeons (squabs) meat is yet to be given recognition. Pigeon farming is at the traditional small scale/backyard level under the extensive (household labour) and semi-intensive management systems in most regions of Africa, Nigeria inclusive (Sule *et al.*, 2024; Husein, Agbolosu and Ishaq, 2023; Kalita and Borah, 2023; NRC, 1991). In these regions the pigeons roam about freely and scavenge around family compounds, feeding on any available farm/harvest residues. The rearer provides the birds with supplements (grains) and the production is usually for family consumption and generation of petty cash / income for the family (Nideou *et al.* 2023; NRC, 1991). Pigeon production has not been given the required attention in comparison to other poultry types (chicken, turkey, guinea fowl, geese etc) even though pigeons have potentials as an alternative animal protein source in Africa. Animal sources of protein are usually expensive and thus most people divert to plant protein which is relatively cheap. Animal protein contains most of the essential amino acids in particular the proteins that are lacking in protein of plant origin namely: methionine, lysine and tryptophan and for this reason animal proteins are described as first class protein, its consumption is highly recommended but due to high cost of animal protein, people decide to switch to plant protein thereby leading to high manifestation of protein deficiency. Micro livestock (rabbit, guinea pig, pigeon, etc) lend themselves to economic riches that are not easily filled by large livestock. Much of their potentials are for subsistence production and they are promising for the many peasants, who are now unable to purchase meat, milk, cheese or eggs due to poor economic situations. These people can afford only animals that can be raised within the homes or backyard under ambient climatic conditions and on feeds that are cheap and easily available. Pigeon production has the potentials for sustainable enhancement of reduction in animal protein deficiency, increased rate of poverty reduction, reduced unemployment and improved socio-economic status of the rural poor community (Asaduzzaman, 2009). Pigeons are kept in almost all parts of Nigeria but not as common as poultry (FDLPCS, 1992). Despite the food value of pigeon for meat only few people engage in it's domestication and its consumption as a source of animal protein in diet in Nigeria (Adawy and Abdel-Wareth, 2023).. In spite of attempts to solve the problem of protein deficiency and malnutrition in Nigeria, it has continuously increased due to the astronomic cost of conventional sources of animal protein (beef, milk, egg. chicken inclusive) which makes them unavailable to the rural populace . This chapter therefore explores the potentials of pigeon meat as an alternative and readily available cheap source of animal protein.

Advantages of Pigeon Farming

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

Pigeon farming has tremendous advantages which include:

- i. Easy management (require little or no handling)
- ii. Require less housing space
- iii. Can be reared easily on roof tops or home yards
- iv. Fast growing and short reproductive cycle
- v. Female starts laying at 5-6months of age at the rate of 2 eggs / month
- vi. They brood the young with little intervention
- vii. Very short hatching period (18days)
- viii. Squabs (baby pigeon) reach table size at 3-4 week of age (thus meat production is sustained compared to other livestock)
- ix. Requires very little investment
- x. Scavenge for food thus cost of feeding is low
- xi. No taboos against pigeon meat
- xii. Meat is highly acceptable and thus in high and constant demand with high premium
- xiii. Entertainment / recreation. People keep them as pets and ornamental birds and one can spend quality time watching their activities during leisure time
- xiv. Source of natural beautification as ornamental birds (their plumages are naturally beautiful)
- xv. Maximum profit with small start-up capital and comparatively low labour cost (feed and housing cost)
- xvi. Provides additional supplemental income
- xvii. Less predisposed to disease compared to other forms of poultry
- xviii. Droppings is a useful bio-fertilizer (manure) for crops and family gardens
- xix. Feathers / plumage used in production of toys
- xx. They keep the environment safe by eating different types of insects
- xxi. Used as laboratory animals in genetic and hormonal studies.
- xxii. Pigeons are extensively used for scientific research notably in physiology and psychology
- xxiii. Homing pigeons are used to carry messages
- xxiv. They are used as pets for sports (racing)
- xxv. The squab is in great demand by sick people and people with nutritional challenges (digestive issues). It is a source of palatable, delicious and easily digestible animal protein (Ghosh *et al.* 2013). The squab contains large proportions of soluble protein and smaller proportions of connective tissues than most forms of meat

Appearance and Size

Pigeons have small head, plump, full breasted bodies, soft and dense plumage. They weigh between 0.5-1kg, however, a few large breeds weigh up to 1.4kg, the size of small domestic chicken.

Habitat and Environment

The domestic pigeon can be raised in temperate and tropical zones including arid and humid regions. However, cold climates do not favour squab production while hot climates promote vermin and diseases.

Distribution

Domestic descendants of the rock pigeon of Europe and Asia are reared today in almost any country (NRC, 1991).

Slaughtering/Dressing

By 28 days, (4 weeks) they should be culled for the table, while they are most tender and before they start to become flighty and lose condition. They are killed and plucked in the same way as other fowls. However, their skins are very tender as such extra care is needed to prevent damage during the dressing process.

Pigeon Meat

There are mainly 300 breeds of pigeon around the world, which are categorized into two types as follows:

1. Meat producing breed (utility)
2. Entertaining/sport/ exhibition/fancy/ornamental breed

Mostly, pigeons are traditionally kept for the purpose of their meat. Baby pigeon meat (squab) is softer and tastier than adult meat. The squabs are harvested just before full feather development and before the youngster has started to fly, usually at 21-30 days of age. At this time the feathers are nearly mature and the meat is soft and tender and their body weight is usually appreciable (the ratio of flesh to inedible parts is highest). Once flying begins, the meat becomes tougher and body weight decreases. The weight of pigeons depends on the breed, nutrition and other factors, but usually ranges from 340-680g (NRC, 1991). Rearers normally expect an average of 12-14 squabs/pair/year. However, this depends on environment and management (NRC, 1991). The birds attain market weight at 450-500g. The market age is usually 4-5weeks (120-150days).

Egg Production

Pigeons are traditionally kept for their meat. They however lay eggs in clutches. Male and female pigeons live in pairs. They collect straws and build a small nest. They start laying at the age of 5-6 months and they lay a pair of eggs every month. Both male and female pigeon hatch the eggs one after the other. It takes 17-18 days to hatch the eggs. Artificial nest can be made for the birds if required. Since the eggs are relatively very small in size, squab production is highly profitable compared to consumption of the eggs

Behaviour of Pigeons

Pigeons mate in pairs and remain as such for life. Once a pair have gone through the courting stage and mated, both are ready to build a nest and hatch young ones together. The mated couples (male and female birds) remain devoted to each other as long as they live or for as long as they are allowed to stay together. However, if mating is disrupted by separation or death, the birds will mate again with other birds/partners. The birds play equal role in nest building, incubation and care for the young. Pigeons typically lay two (2) eggs/clutch/year. A total of 8 clutches/year is not uncommon for a breeding pair. The incubation period is 17-19 days. Unlike most birds, pigeons drink by inserting their beaks into the water and sucking up a continuous draft. Growth and egg production in pigeons declines rapidly after the third year. In domesticated pigeons, sexual maturity (measured as age at first egg) is reached at 120-150 days. Courtship is characterized by cooing, prancing and displays of spread, lowered tail feathers.

Feeding

Naturally pigeons eat insects, leaves, fruits and some invertebrates (NRC, 1991). Generally, scavenger pigeons eat wheat, broken maize, sorghum, paddy rice, legumes, weed seeds and some green vegetables. They also eat spilled grains from feed mills and farm fields. Pigeons require a balanced ration with 15-16% protein to attain proper growth and maximum meat production. The feed intake range of pigeons is 40-50g/pigeon/day. Pigeons also require oyster shell, limestone, salt, grit mixture, bone powder and mineral supplements to attain fast growth. The birds require grit for provision of minerals and to allow the birds to grind the feed in their gizzards. Squabs (baby pigeons) normally do not require extra feed for 5-7 days. They take crop milk (pigeon milk), which is made from their parents stomach for the next 10 days. The female pigeon softens the food in her beak and then feeds the squabs. As soon as the birds are able to fly, they fend and feed for themselves without parental influence or indulgence. The bird must also be given water because they drink water and also like to have a bath in the same water. The water container (pot) should be cleaned on daily basis to avoid disease occurrence.

Brooding and Reproduction

Pigeons are reared in pairs. Most Columbidae species are generally social birds and they mate for life and can become territorial during the breeding season. A pair of male and female birds stays together as couples throughout their life time. They collect straws through mutual effort to build a comfortable nest for them to live in. The female starts laying eggs at 5-6 months of age and lay a pair of eggs every month and both male and female can incubate the eggs, which hatches in 17-18

SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5

days. The eggs are laid in clutches. The female pigeon can lay eggs for up to 5 years of age. The male and female pigeon produce crop milk, which is similar to mammalian milk. The crop of squabs (young pigeons) contains “crop milk”, obtained from their parents, which acts as feed for the first 4 days. In the next 10 days, both male and female pigeon feeds the young birds and thereafter they begin taking feed on their own by 21-30 days. The egg weight is about 20g and the egg shell is white in colour.

Disease

Disease infestation in pigeons is very low compared to any other poultry birds. However, they suffer mainly from tuberculosis (TB), paratyphoid, cholera, pox, newcastle, influenza, etc. Aside from these they can also suffer from malnutrition diseases as well as lice infestation. These diseases are however avoidable with experienced veterinarian consultations and proper management practices as follows:

- Keep the pigeon house clean and germ free
- Isolate disease afflicted birds from healthy birds
- Strict adherence to vaccination regime/schedule
- Keep pigeons free from worms
- Supplementary feed should be given to them to avoid malnutrition diseases

Husbandry

Under extensive conditions where the birds are released each day to feed themselves, almost no land is needed. Under intensive conditions, where the birds spend their lives in confinement, a land of half hectare can be enough space to raise 2,000 pairs. Free ranging pigeons forage over a wide area than most domestic fowls because they fly out to find their feed. Nutrient requirement protein (13.5-15%); carbohydrates (60-70%), fat (2-5%) and fibre (5%) are similar to those of chickens and other fowls (making allowance for the energy needed for flying). Commercial feed and other supplements if needed at all are made available (NRC, 1991). Unlike chickens, pigeons do not prefer communal roosts. Instead they prefer nesting shelves at least two (2) for each breeding pair. The shelves are normally placed in dark corners and fitted with low walls to keep eggs from rolling off. Pigeons are easily trained to recognize “home”. The wing feathers are clipped and the birds are fed close to the dovecote and by the time the wings become unclipped, their homing instincts become developed (NRC, 1991). Alternatively, freshly captured pigeons may be trained by confining them to the dovecote for at least one (1) week. Normally, a little grain is provided in the morning to ensure the birds return home. Thereafter the birds are left to obtain more food for themselves. In captivity, the life span of pigeon species is 10-15 years, however, occasionally, they may live up to 20 years.

Housing

Housing is of paramount importance in pigeon rearing. Most pigeons and doves can be maintained outdoors when the climate is mild or moderate. Any water proof house which is easy to clean is

suitable for rearing pigeons. Many traditional pigeon houses (cotedove) are built of earthenware pots. In Asia and Europe, wooden pigeon towers are also used (NRC, 1991). The following considerations with regards to housing should be put into perspective while managing pigeons:

- A predator (cat, dog, mouse) and rodent – proof flight cage is normally recommended for the birds, thus the pigeon house should be built in a high place.
- Ventilation should be provided in the house, particularly for large flocks (for easy circulation of air)
- Optimum lightening should be ensured (light bulb can be put in place)
- Prevention of entry of rain water inside the house (avoid damp conditions)
- Cleaning of the house at least weekly to prevent the growth of molds and for proper maintenance of hygienic conditions
- Keep house clean and dry always
- Make a door in every room measuring 10x10cm
- Keep food and water constant close to the house
- Keep some straws close to the house so that the birds can make their bed on their own
- Keep water and sand near the house, since they can clean themselves with water and dust
- Build tin wood or tin bamboo packing boxes. Since pigeons and doves spend most of their time on their feet, they require comfortable perching surfaces in form of flat shelves or branch type shelves. These shelves are good for courting, napping etc.
- Each pigeon requires about 30cm x 30cm height and 30cm width space

Research Concerns

Even though, the pigeon production industry remains traditional, it is constantly evolving and deserves scientific research and investigations in the area of diseases, husbandry, feeding and housing for better understanding and harnessing of its potentials towards promoting protein food sufficiency (meat and egg), generation of income, poverty reduction and employment generation amongst resource limited population in rural communities.

Bibliography

- AAV. (2021). Association of Avian Veterinarians. Pigeon Husbandry. Retrieved from www.cdn.ymaws.org/aav/pigeon-husbandry.html.
- Adawy, A.M.G & Abdel-Wareth, A.A.A.(2023).Productive performance and Nutritional of domesticated pigeons. Present status and future concerns. *SVU-International Journal of Agricultural Sciences*, 2023;5(2):160-167.
- Akilu, H. A. (2007). Village Poultry in Ethiopia. Socio-terminal analysis and learning with farmers (unpublished PhD thesis, Wageningen University).
- Almajiri, I. M., Ibrahim, T., Ja'afar, A. M., Shuaibu, A., Umar, H. A. and Ma'aruf, B. S.(2019). A Study on Traditional Pigeon Farming And Breeding Practices In Gamawa L.G.A, Bauchi State .*International Journal of Recent Advances in Multidisciplinary Research*,06(04), 4817-4820
- Almeida da Silva, G., Bernardi, T.L., Carvalho Schaker, P.D., Menegotto, M. and Valente, P. (2012). Rapid yeast DNA extraction by boiling and freeze-thawing without using chemical reagents and DNA purification. *Braz. Arch. Biol. Technol.* 55, 319–327.
- Amal, S., Sahar, A., EL-Rahim Y. A. A. and Aggour, W. (2014).A field study on pigeon production systems governorate, Egypt *Egyptian Poultry Science Journal*, 34(4): 1040-1053.
- Angelier, F., J. C. Wingfield, S. Tartu, and O. Chastel. 2016. Does prolactin mediate parental and life-history decisions in response to environmental conditions in birds? A review. *Horm. Behav.* 77:18– 29.
- Arockia, A. and Indianraj, R.N (2023). Pigeon Bird and its Uses in Agriculture *International Journal of Science and Research (IJSR)*SJIF (2022): 7.942 2.
- Asaduzzaman, M., Mahiuddin, M., Howlider, M. A. R., Hossain, M. M. and Yeasmin, T. (2009). Pigeon farming in Gouripur Upazila of Mymensingh district. *Bangladesh Journal of Animal Science*, 38(1-2):142-150.
- Audu, P. A., Oniye, S. J. and Okechukwu, P. U. (2004). Helminth parasites of domesticated pigeons (*Columba livia domestica*) in Zaria, Nigeria. *Journal of Pest, Disease and Vector Management*, 5: 356-360.
- Aviles-Esquivel, D. F., Martínez, A. M., Landi, V., Álvarez, L. A., Stemmer, A., Gómez-Urviola, N. and Delgado, J. V. (2018). Genetic characterization of South America domestic guinea pig using molecular markers. *Tropical and Subtropical Agroecosystems*, 21(1), 1–10. <https://doi.org/10.56369/tsaes.2588>
- Bament, W. (2012). VN's guide to guinea pigs: Handling, nutrition, nursing. *VN Times*, 12(10), 25–27.

- Balogun, B.I. (2009). Pigeon production for sustainable development. A conference paper, School of Vocational and Technical, Federal College of Education, Zaria. pp1-12.
- Basu, S., Banerjee, D., Chowdhury, R. and Bhattacharya, P. (2018). Controlled release of microencapsulated probiotics in food matrix. *Journal of Food Engineering*, 238, 61–69. <https://doi.org/10.1016/j.jfoodeng.2018.06.005>
- Bislava, M. B., Igwebuikwe, J. U., Buba, S. and Bukar, A. I. (2022). Understanding standard for guinea pig production in Nigeria: A review. *Nigerian Journal of Animal Science*, 24, 90–99.
- Bu, Z. P., Xie, S. Y., Fu, H. B., Tong, and Dai, X. (2015). Effect of energy and protein levels on performance, egg quality, and nutri-ent digestibility of laying pigeons. *J. Appl. Poult. Res.* 24:371–379.
- Burgos-Paz, W., Cerón-Muñoz, M. and Solarte-Portilla, C. (2011). Genetic diversity and population structure of the Guinea pig (*Cavia porcellus*, Rodentia, Caviidae) in Colombia. *Genetics and Molecular Biology*, 34(4), 711–718. <https://doi.org/10.1590/S1415-47572011005000057>
- Caneschi, A., Bardhi, A., Barbarossa, A. and Zaghini, A. (2023). The use of antibiotics and antimicrobial resistance in veterinary medicine, a complex phenomenon: A narrative review. *Antibiotics*, 12(3), 487. <https://doi.org/10.3390/antibiotics12030487>
- Cano, J., Carcelén, F., Ara, M., Quevedo, W., Alvarado, A. and Jiménez, R. (2016). Efecto de la suplementación con una mezcla probiótica sobre el comportamiento productivo de cuyes (*Cavia porcellus*) durante la fase de crecimiento y acabado. *Revista de Investigaciones Veterinarias del Perú*, 27(1), 51. <https://doi.org/10.15381/rivep.v27i1.11458>
- Cárdenas, L. Á. (2018). Características productivas y tecnológicas de la carne de cuy (*Cavia porcellus*) utilizando dietas basadas en pisonay (*Erythrina* sp). *Revista de Investigaciones Altoandinas -Journal of High Andean Research*, 20(4), 451–476. <https://doi.org/10.18271/ria.2018.422>
- Carhuaricra Huaman, D. E., Luna Espinoza, L. R., Rodríguez Cueva, C. L., Duran Gonzales, C. G., Rosadio Alcántara, R. H., Setubal, J. C. and Maturrano Hernández, L. (2022). Genomic characterization of salmonella typhimurium isolated from guinea pigs with salmonellosis in Lima, Peru. *Microorganisms* [Internet], 10(9), 1726. <https://doi.org/10.3390/microorganisms10091726>
- Ciminari, M.E., Mayono, D., Chediack, J.G. and Caviede, V.E. (2005). Feral Pigeons in Urban Environments: Dietary Flexibility and Enzymatic Digestion. *Revista Chilena Historica Natural* 78:267-279
- Chen, M. J., Fu Z. S., Jiang, G., Wang, X. Q., Yan, H. C. and Gao, C. Q. (2020a.) Targeted disruption of TORC1 retards young squab growth by inhibiting

- the synthesis of crop milk protein in breeding pigeon (*Columba livia*). *Poult. Sci.* 99:416–422.
- Chen, M. X., Li, X. G., Yan, H. C., Wang, X. Q. and Gao, C. Q. (2016). Effect of egg weight on composition, embryonic growth, and expression of amino acid transporter genes in yolk sac membranes and small intestines of the domestic pigeon (*Columba livia*). *Poult. Sci.* 95:1425–1432.
- Chen, M. J., Pan, N. X., Wang X. Q., Yan, H. C., and Gao, C. Q. (2020b). Methionine promotes crop milk protein synthesis through the JAK2-STAT5 signaling during lactation of domestic pigeons (*Columba livia*). *Food Funct.* 11:10786–10798.
- Chen, D., Tan H. Z., Liu S. B., Peng Y. Z., Zou, Y., Z. Su, Q., Liu, Y. H. and Li, H. B. (2018). Effects of dietary crude protein level on growth performance, slaughter performance and feather growth in artificial feeding pigeons from 14 to 33 days of age (In Chinese). *Feed Industry* 39:27–30.
- Clinical and Laboratory Standards Institute (CLSI). Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts, 3rd ed.; CLSI Guideline M44; CLSI: Wayne, PA, USA, 2018.
- Pietras-Ozga, D., Ignaciuk, S. and Urban-Chmiel, R. (2021). Biodiversity of *Ligilactobacillus salivarius* strains from poultry and domestic pigeons. *Animals (Basel)* 11:972.
- Didier, M. (2022). Doping in Racing Pigeons (*Columba livia domestica*): A Review and Actual Situation in Belgium, a Leading Country in This Field." *Veterinary Sciences*, 9(2): 42. <http://dx.doi.org/10.3390/vetsci9020042>.
- Dietz, M. W., Salles, J. F., Hsu, B. Y., Dijkstra, C., Groothuis, T. G. G., van der Velde, M., Verkuil, Y. I. and Tieleman, B. I. (2019). Prenatal transfer of gut bacteria in rock pigeon. *Microorganisms* 8:61.
- Ding, H., Lin, Y., Zhang, T., Chen, L., Zhang, G., Wang, J., Xie, K., and Dai, G. (2021). Transcriptome analysis of differentially expressed mRNA related to pigeon muscle development. *Animals (Basel)* 11:2311.
- Ding, J., Liao, N., Zheng, Y., Yang, L., Zhou, H., Xu, K., Han, C., Luo, H., Qin, C., Tang, C., Wei, L., and Meng, H. (2020). The composition and function of pigeon milk microbiota transmitted from parent pigeons to squabs. *Front. Microbiol.* 11:1789.
- FDLPCS (1992). Federal Department of livestock and pest control services. Resource inventory and management limited. Vol. I Exclusive and Atlas, 13.

- FDLPSC (1992) Federal Department of Livestock and pest control services. Resource inventory and management. Nigerian livestock resources survey. Vol. II: national synthesis, pg. 284-287
- FDLPSC (1992) Federal Department of livestock and pest control services. Resource inventory and management limited. Vol. III: state report.
- FDLPSC (1992) Federal Department of livestock and pest control services. Resource inventory and management. Nigerian livestock resources survey. Vol. IV: urban reports and commercially managed livestock survey report.
- Gao, H. J. (2022). Production situation of China pigeon industry in 2021 and trend outlook (in Chinese). *Anim. Agric.* 4:44–46.
- Gao, C. Q., Wang X. H., Hu X. C., Yan, H. C. and Wang, X. Q. (2016a). Effects of dietary crude protein levels on growth performance, carcass characteristics, meat quality of squabs and laying performance of breeding pigeons (in Chinese). *J. South China Agric. Univ.* 37:1–6.
- Gao, C. Q., Yang, J. X., Chen, M. X., Yan, H. C. and Wang, X. Q. (2016b). Growth curves and age-related changes in carcass characteristics, organs, serum parameters, and intestinal transporter gene expression in domestic pigeon (*Columba livia*). *Poult. Sci.* 95:867–877.
- Ge, P., Ma, H., Li, Y., Ni, A., Isa, A. M., Wang, P., Bian, S., Shi, L., Zong, Y., Wang, Y., Jiang, L., Hagos, H., Yuan, J., Sun, Y., and Chen, J. (2021). Identification of microRNA-associated-ceRNA networks regulating crop milk production in pigeon (*Columba livia*). *Genes (Basel)* 12:39.
- Gensollen, T., Iyer, S. S., Kasper, D. L., and Blumberg, R. S. (2016). How colonization by microbiota in early life shapes the immune system. *Science* 352:539–544.
- Gillespie, M. J., Crowley, T. M., Haring, V. R., Wilson, S. L., Harper, J. A., Payne, J. S., Green, D., Monaghan, P., Donald, J. A., Nicholas, K. R., and Moore, R. J. (2013). Transcriptome analysis of pigeon milk production-role of cornification and triglyceride synthesis genes. *BMC Genom.* 14:169.
- Gillespie, M. J., Stanley, D., Chen, H., Donald, J. A., Nicholas, K. R., Moore, R. J. and Crowley, T.M. (2012). Functional similarities between pigeon 'milk' and mammalian milk: induction of immune gene expression and modification of the microbiota. *PLoS One* 7 e48363.
- Giacobbe, D.R., Maraolo, A.E., Simeon, V., Magnè, F., Pace, M.C., Gentile, I., Chiodini, P., Viscoli, C., Sanguinetti, M. and Mikulska, M. (2020) Changes in the relative prevalence

- of candidaemia due to non-albicans *Candida* species in adult in-patients: A systematic review, meta-analysis and meta-regression. *Mycoses*, 63:334-342.
- He, Z. H., Ding, J. T., Shen, M. M., Li, X. R., Zhou, K., and Wu, M. (2012). Analysis of the growth of squab and the nutritional level of crop milk (in Chinese). *Forei. Anim. Husb. Pig Poult.* 32:48–50.
- Hu, X. C., Gao, C. Q., Wang, X. H., Yan, H. C., Chen, Z. S., and Wang, X. Q. (2016). Crop milk protein is synthesised following activation of the IRS1/Akt/TOR signalling pathway in the domestic pigeon (*Columba livia*). *Br. Poult. Sci.* 57:855–862.
- Husein, S.M.A., Agbolosu, A.A.& Ishaq, N.(2023). Characterization of Pigeon (*Columba livia domestica*) Production Systems and Marketing in Northern Ghana. *The Pharma Innovation Journal*, SP-12(12): 2411-2413
- Ibrahiem, L. (2013). Structural changes in pigeon crop mucosa during crop milk secretion. *Zag. Vet. J.* 41:113–123.
- Janssens, G. P., Hesta, M., Debal, V., Debraekeleer, J. and de Wilde R. O. (2000a). L-carnitine supplementation in breeding pigeons: impact on zootechnical performance and carnitine metabolism. *Reprod. Nutr. Dev.* 40:535–548.
- Janssens, G. P. J., Hesta, M., Debal, V., and de Wilde, R. O. M. (2000). The effect of feed enzymes on nutrient and energy retention in young racing pigeons. *Ann. Zootech.* 49:151–156.
- Ji, F., Zhang, S., An, Y., Wang, Z., Shao, Y., Du, S., Xing, L., and Sun, X. (2022). Influence of dietary phosphorus concentrations on the performance of rearing pigeons (*Columba livia*), and bone properties of squabs. *Poult. Sci.* 101:101744.
- Ji, F., Zhang, D., Shao, Y., Yu, X., Liu, X., Shan, D., and Wang, Z. (2020). Changes in the diversity and composition of gut microbiota in pigeon squabs infected with *Trichomonas gallinae*. *Sci. Rep.* 10:19978.
- Jiang, S. G., Pan, N. X., Chen, M. J., Wang, X. Q., Yan, H. C., and Gao, C. Q. (2019). Effects of dietary supplementation with DL-methionine and DL-methionyl-dl-methionine in breeding pigeons on the carcass characteristics, meat quality and antioxidant activity of squabs. *Antioxidants (Basel)* 8:435.
- Jin, C. L., Ye, M., Song, Z. W., Zhang, Z. M., Gao, C. Q., Yan, H. C., and Wang, X. Q. (2022). Lysine interacts with *frizzled7* to activate β -catenin in satellite cell-participated skeletal muscle growth. *J. Agric. Food Chem.* 70:3745–3756.

- Jin, C. L., Zhang, Z. M., Song, Z. W., Gao, C. Q., Yan, H. C., and Wang, X. Q. (2020). mTORC1-mediated satellite cell differentiation is required for lysine-induced skeletal muscle growth. *J. Agric. Food Chem.* 68:4884–4892.
- Jorgensen, J. M., Young, R., Ashorn, P., Ashorn, U., Chaima, D., Davis, J. C. C., Goonatileke, E., Kumwenda, C., Lebrilla, C. B., Maleta, K., Sadalaki, J., Totten, S. M., Wu, L. D., Zivkovic, A. M., and Dewey, K. G. (2021). Associations of human milk oligosaccharides and bioactive proteins with infant morbidity and inflammation in malawian mother-infant dyads. *Curr. Dev. Nutr.* 5nab072.
- Jin, C. L., Ye, M., Song, Z. W., Zhang, Z. M., Gao, C. Q., Yan, H. C. and Wang, X. Q. (2022). Lysine interacts with frizzled7 to activate b-catenin in satellite cell-participated skeletal muscle growth. *J. Agric. Food Chem.* 70:3745–3756.
- Jin, C. L., Zhang, Z. M., Song, Z. W., Gao, C. Q., Yan, H. C., and Wang, X. Q. (2020). mTORC1-mediated satellite cell differentiation is required for lysine-induced skeletal muscle growth. *J. Agric. Food Chem.* 68:4884–4892.
- Jorgensen, J. M., Young, R., Ashorn, P., Ashorn, U., Chaima, D., Davis, J. C. C., Goonatileke, E., Kumwenda, C., Lebrilla, C. B., Maleta, K., Sadalaki, J., Totten, S.M., Wu, L. D., Zivkovic, A. M., and Dewey, K. G. (2021). Associations of human milk oligosaccharides and bioactive proteins with infant morbidity and inflammation in malawian mother-infant dyads. *Curr. Dev. Nutr.* 5nab072.
- Kalita, J.J. and Borah, D.(2023). Comprehensive review on pigeon farming: A sustainable livelihood option for rural communities in Assam .*Research & Reviews: A Journal of Veterinary Science and Technology*, 12(3):35-42
- Liang, X., Yu, J., Wang, H., and Zhang, Z. (2018). Post-hatching growth of the pectoralis muscle in pigeon and its functional implications. *Anat. Rec. (Hoboken)* 301:1564–1569.
- Ma, Y., Feng, S., Wang, X., Qazi, I. H., Long, K., Luo, Y., Li, G., Ning, C., Wang, Y., Hu, S., Xiao, J., Li, X., Lan, D., Hu, Y., Tang, Q., Ma, J., Jin, L., Jiang, A. and Li, M. (2018). Exploration of exosomal micro-RNA expression profiles in pigeon 'Milk' during the lactation period. *BMC Genom.* 19:828.
- Ma, H., Ni, A., Ge, P., Li, Y., Shi, L., Wang, P., Jing. Fan., Isa A. M., Sun, Y., and Chen, J. (2020). Analysis of long non-coding RNAs and mRNAs associated with lactation in the crop of pigeons (*Columba livia*). *Genes (Basel)* 11:201.

- Mabuchi, Y., and Frankel, T. L. (2016). Functions of innate and acquired immune system are reduced in domestic pigeons (*Columba livia domestica*) given a low protein diet. *R. Soc. Open Sci.* 3:150408.
- Mahdy, M. A. A. (2021). Comparative morphological study of the oro-pharyngeal floor of squabs and adult domestic pigeons (*Columba livia domestica*). *Microsc. Res. Tech.* 84: 99–511.
- Meleg, I., Dublec, K., Vincze, L. and Horn, P. (1999). Effect of dietary crude protein level on reproductive traits of commercial pigeons in different production terms. *Acta Agraria. Kaposvariensis* 2:247-253.
- Mohamed, R. A., Shukry, M., Mousa-Balabel, T. M., and Elbassiouny, A. A. (2016). Assessment of plasma prolactin and nest defense behaviour during breeding cycle of pigeon (*Columba livia domestica*). *J. Environ. Agric. Sci.* 7:19–22.
- Moazama, Batool, (n.d). Pigeon Farming Poultry Production. *BioScientific Review*, 6(4): 27
- NRC: National Research Council (1991). Micro livestock: little known small animals with a promising economic future. Pigeon. National academy press, Washington, D.C. pg 136-145 small holders guide. University services limited. Yaba lagos.
- Nideou, D.M, Tellah, Adoum, I.Y, Borso, A.J. Onagbessan, O., Kokou, T. and Logtene, Y.M. (2023). Morphological and Zootechnical Characteristics of the Local Chicken in the Urban Area of Abeche City, *International Journal of Poultry Sciences* 22:5-11
- Nideou, D., Tellah, M., Leng T., Djasrangue, S. and Mopate, L.Y. (2023). Breeding Practice and Productivity of Domestic Pigeon (*Columbia livia domestica*) in the Urban Area of Abeche City. *Journal of Applied Biosciences* 188:19813-19822
- Ley Orgánica de Sanidad Agropecuaria. Asamblea Nacional, República del Ecuador. Available online:https://www.gob.ec/sites/default/files/regulations/2018-09/Documento_Ley%20Org%C3%A1nica%20de%20Sanidad%20Agropecuaria.pdf
- Gardes, M. and Bruns, T.D. (1993). ITS primers with enhanced specificity for Basidiomycetes: Application to the identification of mycorrhizae and rusts. *Mol. Ecol.* 2, 113–118.
- Ghosht, K.K., Sikder, S., Chowdhury, S., Ray, M.K., Sarker, A, and Chanda, G.C. (2013). A Study on Management System of an Intensive Pigeon Farming at Chittagong. *Wayamba Journal of Animal Science* pp: 623-628
- Jin, C., He, Y., Jiang, S., Wang, X., Yan, H., Tan, H and Gao, C. (2023). Chemical Composition of Pigeon Crop Milk and Factors affecting its Production: A Review *Poultry Science* pp14
- Kabir, M.A. (2018). Necessary Steps to Establish a First-Time Pigeon Farm. *Journal of Dairy, Veterinary and Animal Research* 7(6): 248-251

- Papon, N.; Savini, V.; Lanoue, A.; Simkin, A.J.; Crèche, J.; Giglioli-Guivarc'h, N.; Clastre, M.; Courdavault, V.; Sibirny, A.A. and Candida, G. (2013). Biotechnological applications, perspectives for biological control, emerging clinical importance and recent advances in genetics. *Curr. Genet.* 2013, 59, 73–90.
- Pigeon Husbandry *Association of Avian Veterinarians* (AAV)
- Ramachandran, A. V., and Patel, M. M. (1989). Seasonal differences in glucose tolerance and insulin response of pinealectomized pigeons (*Columba livia*). *J. Pineal. Res.* 6:209–219.
- Sales, J. and Janssens. G. P. J. (2003a). Methods to determine metabolizable energy and digestibility of feed ingredients in the domestic pigeon (*Columba livia domestica*). *Poult. Sci.* 82:1457–1461.
- Sales, J., and Janssens. G. P. J. (2003b). Nutrition of the domestic pigeon (*Columba livia domestica*). *World Poult. Sci. J.* 59:221–232.
- Shao, Y., Ma, W., Ji, F., Sun, X., Du, S., Li, X., Li, Q. and Wang, Z. (2021). Exploration of proteomics analysis of crop milk in pigeons (*Columba livia*) during the lactation period. *ACS Omega* 6:27726–27736.
- Shetty, S., Bharathi, L., Shenoy, K. B., and Hegde, S. N. (1992a). Bio-chemical properties of pigeon milk and its effect on growth. *J. Comp. Physiol. B* 162:632–636.
- Shetty, S., and Hegde, S. N. (1991). Changes in lipids of pigeon "milk" in the first week secretion. *Lipids* 26:930–933.
- Shetty, S., and Hegde, S. N. (1993). Pigeon milk: a new source of growth factor. *Experientia* 49:925–928.
- Shetty, S., Hegde, S. N., and Bharathi, L. (1992b). Purification of a growth factor from pigeon milk. *Biochim. Biophys. Acta* 1117:193–198.
- Sossinka, R. (1982). Domestication in birds. *Avian Biol.* 6:373–403. Stadie, W. C., N. Haugaard, and M. Perlmutter. 1948. The effect of insulin upon pyruvate utilization by pigeon muscle. *J. Biol. Chem.* 172:567–577.
- Skinner, J. (2004). *Animal Sciences: Pigeons*, Winsconsin 4-H Publications. University of Winsconsin, pp. 12
- Sule, K., Adeleke, M.A., Adegbenro, M., Folorunso, O.R and Onibi, G.E. (2024). Management and challenges of pigeon production in South-West Nigeria. *International Journal of Novel Research in Life Sciences*, 11(1): 14 – 24.

- Sule, K., Adeleke, M.L., Adegbenro, M., Folorunso, O.R., and Onibi, G.E. (2024). People's Perception of Domestic Pigeon (*Columba livia*) in South-West, Nigeria. *Continental J. Applied Science*, 19(1), 64–75. <https://doi.org/10.5281/zenodo.10706902>
- Vandeputte-Poma, J. (1980). Feeding, growth and metabolism of the pigeon, *Columba livia domestica*: duration and role of crop milk feeding. *J. Comp. Physiol.* 135:97–99.
- Waldie, G. A., Olomu, J. M., Cheng, K. M., and Sim, J. (1991). Effects of two feeding systems, two protein levels, and different dietary energy sources and levels on performance of squabbing pigeons. *Poult. Sci.* 70:1206–1212.
- Wan, X. P., Xie, P., Bu, Z., Zou, X. T., and Gong, D. Q. (2019). Prolactin induces lipid synthesis of organ-cultured pigeon crops. *Poult. Sci* 98:1842–1853.
- Wang, F., Li, F. H., Zhan, S. J., Chen, Y., Zhu, X. J. and Guo, K. J. (2018). Study on the energy and protein requirements of meat pigeon under “2+3” feeding pattern (in Chinese). *J. Beijing Univ. Agric.* 33:58–63.
- Wang, X. Q., Li, S. B., Zhan, X., Shu, G. S., Liu, H., Rao, C. X., and Xu, H. J. (2009a). The energy of the breeding pigeons needs to be in the “2+4” production mode of the pigeon breeding (in Chinese). *Cereal Feed Ind.* 4:45–46.
- Wang, X. Q., Li, S. B., Zhan, X., Shu, G., Rao, C. X., and Liu, S. H. (2009b). The crude protein requirement of the breeding pigeon in the “2+4” production mode of the pigeon breeding (in Chinese). *Feed Ind.* 30:59–60.
- Wen, J., Zhao, W., Li, J., Hu, C., Zou, X. and Dong, X. (2022). Dietary supplementation of chitosan oligosaccharide-clostridium butyricum synbiotic relieved early-weaned stress by improving intestinal health on pigeon squabs (*Columba livia*). *Front. Immunol.*13:926162.
- White, T.J.; Bruns, T.; Lee, S. and Taylor, J.(1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In PCR Protocols a Guide to Methods and Applications; Academic Press: San Diego, CA, USA, 1990; pp. 315–322
- Cheng, X. C. (2014). Effects of zinc-methionine on growth performance, intestinal flora and immune function in pigeon squabs. *Br. Poult. Sci.* 55:403–408.
- Xie, W. Y., Fu, Z., Pan, N. X., Yan, H. C., Wang, X. Q. and Gao, C. Q. (2019). Leucine promotes the growth of squabs by increasing crop milk protein synthesis through the TOR signaling pathway in the domestic pigeon (*Columba livia*). *Poult. Sci.* 98:5514–5524.
- Xie, P., Han, M. X., Chen, W. X., Wan, X. P., Xu, Y. G. and Gong, D. Q. (2020). The profiling of amino acids in crop milk and plasma and mRNA abundance of
- SAEREM BOOK CHAPTERS2023: First published 2024: ISBN 978-978-60709-9-5**

- amino acid transporters and enzymes related to amino acid synthesis in the crop tissue of male and female pigeons during incubation and chick-rearing periods. *Poult. Sci.* 99:1628–1642.
- Xie, P., Jiang, X.Y., Bu, Z., Fu, S. Y., Zhang, S. Y. and Tang, Q. P. (2016). Free choice feeding of whole grains in meat-type pigeons: 1. Effect on performance, carcass traits and organ development. *Br. Poult. Sci.* 57:699–706.
- Xie, P., Wang, Y., Wang, C., Yuan, C., and Zou, X. (2013). Effect of different fat sources in parental diets on growth performance, villus morphology, digestive enzymes and colorectal microbiota in pigeon squabs. *Arch. Anim. Nutr.* 67:147–160.
- Xie, P., J. Zhu, G., Liu, Y., Liu, T. W., Xu, Y. G. and Gong, D. Q. (2021). Effect of Akt activation on apoptosis-related gene expression in the crop tissues of male and female pigeons (*Columba livia*). *Poult. Sci.* 100:101392.
- Xu, Q., Jian, H., Zhao, W., Li, J., Zou, X. and Dong, X. (2022a). Early weaning stress induces intestinal microbiota disturbance, mucosal barrier dysfunction and inflammation response activation in pigeon squabs. *Front. Microbiol.* 13:877866.
- Xu, Q. Q., Ma, X. W., Dong, X. Y., Tao, Z. R., Lu, L. Z., and Zou, X. T. (2020). Effects of parental dietary linoleic acid on growth performance, antioxidant capacity, and lipid metabolism in domestic pigeons (*Columba livia*). *Poult. Sci.* 99:1471–1482.
- Xu, Q., Wen, J., Wang, X., Zou, X. and Dong, X. (2021). Maternal dietary linoleic acid altered intestinal barrier function in domestic pigeons (*Columba livia*). *Br. J. Nutr.* 126:1003–1016.
- Xu, Q., Zhao, W., Li, Y., Zou X., and Dong, X. (2022b). Intestinal immune development is accompanied by temporal deviation in microbiota composition of newly hatched pigeon squabs. *Microbiol. Spectr.* 10:e0189221.
- Yang, M. C., and Vohra, P. (1987). Protein and metabolizable energy requirements of hand-fed squabs from hatching to 28 days of age. *Poult. Sci.* 66:2017–2023.
- Yuan, J., Ni, A., Li, Y., Bian, S., Liu, Y., Wang, P., Shi, L., Isa, A. M., Ge, P., Sun, Y., Ma, H., and Chen, J. (2021). Transcriptome analysis revealed potential mechanisms of resistance to trichomoniasis gallinae infection in pigeon (*Columba livia*). *Front. Vet. Sci.* 8:672270.
- Zeng, Q. F., Zhang, B., and Zhang, Z. W. (2003). Study on crop milk (in Chinese). *China Poult.* 21:1–3.

- Zhang, X. Y., Dong, X. Y., Bu, X. C., and Zou, X. T. (2016). Bioactive constituents in pigeon milk during 2-10 days secretion (In Chinese). *Nutr. Feedstuffs* 52:39–42.
- Zhao, W., Liu, Q., Jiang, H., Zheng, M., Qian, M., Zeng, X., and Bai, W. (2022). Monitoring the variations in physicochemical characteristics of squab meat during the braising cooking process. *Food Sci. Nutr.* 10:2727–2735.
- Zhong, C., Tong, D. Q., Zhang, Y. R., Wang, X. Q., Yan, H. C., Tan, H. Z., and Gao, C. Q. (2022). DL-methionine and DL-methionyl-DL-methionine increase intestinal development and activate Wnt/b-catenin signaling activity in domestic pigeons (*Columba livia*). *Poult. Sci.* 101:101644.
- Zhu, J. G., Xie, P., Song, C., Liu, T. W. and Gong D. Q. (2022). Differential expression of glucose metabolism-related genes and AMP-activated protein kinases in crop tissue of male and female pigeons (*Columba livia domestica*) during the incubation and chick-rearing periods. *J. Anim. Physiol. Anim. Nutr. (Berl.)* 107:680–690.

